



Dijet production in diffractive deep inelastic scattering using VFPS at HERA

Julie Delvax


I.I.H.E., Université Libre de Bruxelles

On behalf of the H1 collaboration



XIX International Workshop on Deep-Inelastic Scattering and Related
Subjects

Inclusive Diffraction in H1



Q^2 = virtuality of the γ

x_{IP} = momentum fraction of the p carried by the \mathbb{P}

β = momentum fraction of the \mathbb{P} interacting with the γ


t = momentum squared of the \mathbb{P}

y = inelasticity


$$\frac{d^3 \sigma(x_{IP}, Q^2, \beta)}{dx_{IP} dQ^2 d\beta} = \frac{4\pi\alpha^2}{\beta Q^4} (1 - y + \frac{y^2}{2}) F_2^{D(3)}(x_{IP}, Q^2, \beta)$$

H1 PRELIMINARY

- H1 VFPS Preliminary
- H1 FPS
- H1 LRG Preliminary x 0.81
- H1 LRG Published x 0.81
- H1 2006 DPDF Fit B x 0.81
- - - H1 2006 DPDF Fit B x 0.81 (extrapol.)



Jets in diffraction




No direct access to gluon component



Large uncertainty on gluon distribution


Jets in diffraction



No direct access to gluon component



Large uncertainty on gluon distribution



$z_{\mathbb{P}}$ = fraction of momentum of \mathbb{P} entering in the hard interaction.



Jets via BOSON-GLUON FUSION





Directly sensitive to gluon densities in \mathbb{P}

Jets in DIS

DPDFs to predict jet production via Factorization theorem



Dijet included in the fit \Rightarrow improve accuracy at high β




The Very Forward Proton Spectrometer




Purpose : direct measurement of the scattered proton with high acceptance and low background

Since 2004, stable data since 2006

Located at 220 meters from the interaction point



- 2 stations each with 2 scintillating fibers detector
- 4 planes of 4 trigger tiles




The VFPS

In a simple approximation ($t = 0 \text{ GeV}^2$) position in x of p -impact is proportional to $x_{\mathbb{P}}$

$x_{\mathbb{P}}$ is reconstructed directly from VFPS



Check calibration with exclusive events (ρ) : $e p \rightarrow e p \rho$

Compare $x_{\mathbb{P}}^\rho$ reconstructed from ρ decay tracks and scattered electron to $x_{\mathbb{P}}^{\text{VFPS}}$ reconstructed using VFPS




The VFPS

- High Acceptance $\approx 90\%$ for $0.01 < x_{IP} < 0.025$



- High track reconstruction efficiency $\approx 96\%$
- Low background (Beam Gas + DIS) $< 2\%$

$$\left\{ \begin{array}{l} pA \rightarrow p(A') \\ ep \rightarrow eX \end{array} \right.$$



Selection of events

DIS Cuts

- Electron in SpaCal Calorimeter
- At least one track in VFPS1 OR in VFPS2

Jet cuts

- At least 2 central jets
- Jets selected in $\gamma^* - p$ frame
- $P_{t,jet1}^* > 5.5 GeV$
- $P_{t,jet2}^* > 4 GeV$

Kinematic Range

$$\begin{aligned}0.009 < x_{\mathbb{P}} < 0.024 \\ 5 < Q^2 < 80 \text{ } GeV^2 \\ 0.1 < y < 0.65 \\ -3 < \eta_{j1,j2}^* < 0\end{aligned}$$

Kinematic Reconstruction

$$Q^2 = 4E_{e'}E_e \cos^2\left(\frac{\theta_{e'}}{2}\right)$$

$$y = 1 - \frac{E_{e'}}{E_e} \sin^2\left(\frac{\theta_{e'}}{2}\right)$$

$$x_{\mathbb{P}} = x_{\mathbb{P}}^{VFPs}$$

$$z_{\mathbb{P}} = \frac{x}{x_{\mathbb{P}}} \frac{Q^2 + M_{12}^2}{Q^2} \quad M_{12}^2 = (p_{j1} + p_{j2})^2$$

Jets in DIS with VFPS

Data Sample

2006-2007 $e^+ - e^-$: Lumi= 127.1 pb^{-1} $\Rightarrow 1400$ events selected

MC Sample

Rapgap 31 H1 DPDF 2006 Fit B

NLO Predictions

NLO Jets ++ using DPDF 2007 Jets (x0.83 to account for p -dissociation)

$$\mu_{r,f}^2 : 0.5 * (P_{t,jet1}^2 + P_{t,jet2}^2) + Q^2$$

Scale ($\mu_{r,f}^2 \times 0.25$ and $\times 4$) \oplus Hadronization uncertainties




Cross Section Measurement

Cross Section corrected to the Born Level

Data shown with uncorrelated systematic \oplus statistic uncertainties


Correlated uncertainty of 5% not shown

Jet cross section in DIS with VFPS





Within uncertainties
agreement between
NLO predictions using Fit Jets
2007 and Data
in shape and normalisation



Jet cross section in DIS with VFPS



Good description of the transverse momentum of jets \Rightarrow
Good description of the hard interaction



Jet cross section in DIS with VFPS



Extrapolated DPDF for $z_{IP} > 0.9$

Higher z_{IP} bin in agreement with Fit Jets 2007

Total Cross Section

H1 VFPS Preliminary : 25.3 ± 1.4 (stat.) ± 6.5 (syst) pb

NLO DPDF Fit Jet 2007 : $19.9^{+7.4}_{-4.4} \pm 0.5$ (had.) pb

Conclusions

For the first time measurement of Dijets cross section is made with the VFPS

- HERA II data
- Based on direct detection of the scattered proton
- High acceptance detector \Rightarrow High statistic

Results are in agreement with NLO predictions using DPDF 2007 Fit Jets and confirm validity of factorization theorem in DIS

Possibility of extracting new DPDF in the future